

Post-Op Recovery Guide - (Cholecystectomy)

Diet

- **Start Slow:** Let your appetite guide you; do not force food if you aren't hungry.
- **Hydration:** Drink plenty of liquids.
- **Nausea:** Stick to clear liquids if you feel nauseous, then move to easily digestible foods.
- **Fatty Foods:** Avoid fatty foods for the first few weeks as they may cause discomfort or diarrhea.

Activity

- **Movement:** Walk frequently (at least 5 times daily) to help healing and prevent blood clots.
- **Rest:** Take it easy, but do not stay in bed all day. Change positions regularly.
- **Stairs:** You may walk and use stairs as usual.
- **Lifting/Exercise:** Avoid heavy lifting or vigorous exercise for 4–6 weeks.
- **Driving:** Do not drive for 48–72 hours or while taking narcotic pain medication.

Pain Control

- **Medication:** Take prescribed medications as instructed.
- **Pain Meds:** Do not take pain medicine on an empty stomach.
- **Ice:** Use cold packs 3–4 times daily for 20 minutes during the first 2 days to help with swelling.

Wound Care & Showering

- **Showering:** You may shower within 24 hours after surgery. Let water run over incisions; do not scrub.
- **No Soaking:** Do not use baths, jacuzzis, or pools until cleared.
- **Dressings:** Do not peel off surgical glue (Dermabond) or white tapes (Steri-strips); they will fall off on their own.
- **Bruising:** It is normal and harmless for the surgical site to turn black and blue.

When to Call for Help

Call the Clinic (661) 219-2643 if you have:

- Fever over 100.4°F.
- Drainage or increasing redness around incisions.
- Worsening pain despite medication.

Call 911 or go to the ER for:

- Chest pain or shortness of breath.
- Severe leg swelling.
- Loss of consciousness.
- Severe pain or inability to keep liquids down.

Follow-Up: If you don't have an appointment, call the office within 1–2 days at **661-219-2643**.